

# Seismicity of the Earth 1900–2010

## Japan and Vicinity

Compiled by Susan Rhea, Arthur C. Tarr, Gavin Hayes, Antonio Villaseñor\*, Richard L. Dart, and Harley Benz

\*Institute of Earth Sciences, CSIC, Lluís Solé i Sabarís s/n, 08028 Barcelona, Spain

### TECTONIC SUMMARY

This map shows details of Japan and vicinity not visible in an earlier publication (Tarr and others, 2010). Japan and its island possessions lie across four major tectonic plates: Pacific plate; North America plate; Eurasia plate; and Philippine Sea plate. The Pacific plate is subducted into the mantle, beneath Hokkaido and northern Honshu, along the eastern margin of the Okhotsk microplate, a proposed subdivision of the North America plate (Bird, 2003). Further south, the Pacific plate is subducted beneath volcanic islands along the eastern margin of the Philippine Sea plate. This 2,200 km-long zone of subduction of the Pacific plate is responsible for the creation of the deep offshore Ogasawara and Japan trenches as well as parallel chains of islands and volcanoes, typical of Circumpacific island arcs. Similarly, the Philippine Sea plate is itself subducting under the Eurasia plate along a zone, extending from Taiwan to southern Honshu, that comprises the Ryukyu Islands and the Nansei-Shoto trench.

Subduction zones at the Japanese island arcs are geologically complex and produce numerous earthquakes from multiple sources. Deformation of the overriding plates generates shallow crustal earthquakes, whereas slip at the interface of the plates generates interplate earthquakes that extend from near the base of the trench to depths of 40 to 60 km. At greater depths, Japanese arc earthquakes occur within the subducting Pacific and Philippine Sea plates and can reach depths of nearly 700 km. Since 1900, two great earthquakes occurred off Japan and three north of Hokkaido. They are the M8.4 1933 Sanriku-oki earthquake (Kawakatsu and Seno, 1983), the M8.3 2003 Tokachi-oki earthquake (Miyazaki and others, 2004), M8.4 1958 Etorofu earthquake (Fukao and Furumoto, 1979), the M8.5 1963 Kuril earthquake (Beck and Ruff, 1987), and the M8.3 1994 Shikotan earthquake (Kikuchi and Kanamori, 1995).

Several relevant tectonic elements, plate boundaries and active volcanoes, provide a context for the seismicity presented on the main map. The plate boundaries (Bird, 2003) are known most accurately along the axis of the trenches and are more diffuse or speculative in the Sea of Japan, China, and Russia. The active volcanic arcs (Siebert and Simkin, 2002) follow the Izu, Volcano, and Ryukyu island chains and the main Japanese islands parallel to the Japan trench.

### DATA SOURCES

The earthquakes portrayed on the main map and the depth profiles are taken from two sources: (a) the Centennial earthquake catalog (Engdahl and Villaseñor, 2002) and annual supplements for the interval 1900–2007, where the magnitude floor is 5.5 globally, and (b) a catalog of earthquakes having high-quality depth determinations for the period 1964–2002 and a magnitude range of 5.0 ≤ M ≤ 8.4 (Engdahl, personal comm. 2003).

The nucleation points of great earthquakes (M ≥ 8.3) are designated with a label showing the year of occurrence. Their rupture areas are shown as pale reddish polygons. Major earthquakes (7.5 ≤ M ≤ 8.2) are labeled with the year of occurrence, while earthquakes (8.0 ≤ M ≤ 8.2) are labeled with the year of occurrence and also denoted by a white outline.

The Seismic Hazard and Relative Plate Motion panel displays the generalized seismic hazard of the region (Giardini and others, 1999) and representative relative plate motion vectors using the NUVEL-1A model (DeMets, et al., 1994).


Pre-instrumental seismicity was obtained from the NOAA National Geophysical Data Center (2010) database of significant earthquakes; locations are approximate, based on macro-seismic reports and field investigations. We selected earthquakes with associated reports of moderate to major damage, deaths, an estimated magnitude of 7.5 or greater (if known), or tsunami generation.

Base map data sources include GEBCO 2008, Volcanoes of the World dataset (Siebert and Simkin, 2002), plate boundaries (Bird, 2003), Digital Chart of the World, and ESRI (2002). Slab contours are from Hayes and

### REFERENCES

- Beck, S.L., and Ruff, L.J., 1987, Rupture process of the great 1963 Kuril Islands earthquake sequence—Asperity interaction and multiple event rupture. *Journal of Geophysical Research*, v. 92, no. B13, p. 14123–14138.
- Bird, Peter, 2003, An updated digital model of plate boundaries: *Geochemistry Geophysics Geosystems*, v. 4, no. 3, 52 p.
- DeMets, C., Gordon, R.G., Argus, D.F., and Stein, S., 1994, Effects of recent revisions to the geomagnetic time scale on estimates of current plate motions: *Geophysical Research Letters*, v. 21, p. 2191–2194.
- Engdahl, E.R., and Villaseñor, Antonio, 2002, Global seismicity 1900–1999, in Lee, W.H.K., Jennings, P., Kisslinger, C., and Kanamori, H., eds, *International Handbook of Earthquake and Engineering Seismology*, v. 81(A), chap. 41, p. 1–26.
- ESRI, 2002, *ESRI Data and Maps: ESRI*, Redlands, CA.
- Fukao, Yoshio, and Furumoto, Muneyoshi, 1979, Stress drops, wave spectra and recurrence intervals of great earthquakes—Implications of the Etorofu earthquake of 1958 November 6: *Geophysical Journal International*, v. 57, p. 23–40.
- GEBCO, 2008, The GEBCO\_08\_Grid, ver. 20091120, accessed January 8, 2010 at <http://www.gebco.net/>.
- Giardini, D., Grünthal, G., Shedlock, K., Zhang, P., and Global Seismic Hazards Program, 1999, *Global Seismic Hazards Map*, accessed January 9, 2007 at <http://www.seismo.ethz.ch/GSHAP>.
- Hayes, G., and Wald, D., 2010, Slab models for subduction zones: U.S. Geological Survey Earthquake Hazards Program, accessed February 23, 2010 at <http://earthquake.usgs.gov/research/data/slab/>.
- Kawakatsu, H. and Seno, Tetsuo, 1983, Triple seismic zone and the regional variation of seismicity along the northern Honshu: *Journal of Geophysical Research*, v. 88, no. B5, p. 4215–4230.
- Kikuchi, Matsuyuki, and Kanamori, Hiroo, 1995, The Shikotan earthquake of October 4, 1994—Lithospheric earthquake: *Geophysical Research Letters*, v. 22, no. 9, p. 1025–1028.
- Miyazaki, Shin-ichi, Segall, Paul, Fukuda, Junichi, and Kato, Teryuki, 2004, Space time distribution of afterslip following the 2003 Tokachi-oki earthquake—Implications for variations in fault zone frictional properties: *Geophysical Research Letters*, v. 31, no. L06623, p. 1–4.
- NOAA National Geophysical Data Center, 2010, U.S. Dept of Commerce, accessed March 31, 2010 at <http://www.ngdc.noaa.gov/hazards>.
- Siebert, L., and Simkin, T., 2002, *Volcanoes of the world—An illustrated catalog of Holocene volcanoes and their eruptions*: Smithsonian Institution, Global Volcanism Program Digital Information series, GVP-3, <http://www.volcano.si.edu/world/>, last accessed January 9, 2007.
- Tarr, A.C., Villaseñor, Antonio, Furlong, K.P., Rhea, Susan, and Benz, H.M., 2010, *Seismicity of the Earth 1900–2007: U.S. Geological Survey Scientific Investigations Map 3064*, scale 1:25,000,000.


### PRE-INSTRUMENTAL SEISMICITY 1500–1899



### FIGURE EXPLANATION



### SEISMIC HAZARD AND RELATIVE PLATE MOTION




Any use of trade names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Although this information product, for the most part, is in the public domain, it also contains copyrighted materials as noted in the text. Permission to reproduce copyrighted items for other than personal use must be secured from the copyright owner.






This map was produced on request, directly from digital files, on an electronic plotter.

A PDF for this map is available at <http://pubs.usgs.gov/of/2011/1083-D/>

Suggested citation:  
Rhea, Susan, Tarr, A.C., Hayes, Gavin, Villaseñor, Antonio, Richard L. Dart, and Benz, H.M., 2011, *Seismicity of the Earth 1900–2010, Japan and vicinity*: U.S. Geological Survey Open-File Report 2011–1083-D, scale 1:6,000,000.




### MAP EXPLANATION



### DEPTH PROFILE EXPLANATION

Profiles of earthquake and volcano locations are constructed from the mapped data. Locations of the profile intersection with the surface are drawn in the map and labeled to coincide with the profile label. Box defines extent of earthquakes included in profile. Length of the profile graphic is the same as in the map. Distance in kilometers from the trench axis is indicated in the X direction, depth in kilometers is indicated in the Y direction. There is no vertical exaggeration. See Explanation at side for color key. Not all earth layers, earthquake depths or magnitudes, are visible on every map.



Albers Equal Area Conic Projection

Digital map database and cartography by Susan Rhea and Arthur Tarr

Manuscript approved for publication Month Day, 2011

Albers Equal Area Conic Projection

SCALE 1:6 000 000

0 100 200 400 600 800 Kilometers

